

Starting your patients on **CYSTADROPS**[®] (cysteamine ophthalmic solution) 0.37%


Here are some tips you may want to consider when starting your patients on **CYSTADROPS**:

- 1 Explain how **CYSTADROPS** works
- 2 Review potential results
- 3 Assist with dosing and instillation
- 4 Discuss most common side effects

1. Explain how **CYSTADROPS** works

CYSTADROPS works by:

- 💧 Coating the entire surface of the cornea
- 💧 Penetrating throughout the cornea to reach all layers of the cornea
- 💧 Dissolving cystine crystals in all layers of the cornea



2. Review potential results

Studies of **CYSTADROPS** have shown:

- 💧 A significant reduction in corneal cystine crystals after 90 days of 4x/day instillation
- 💧 The crystal reduction was maintained over time with continuous use

Indications and Usage

CYSTADROPS (cysteamine ophthalmic solution) 0.37% is a cystine-depleting agent indicated for the treatment of corneal cystine crystal deposits in adults and children with cystinosis.

Important Safety Information

- To minimize the risk of contamination, do not touch the dropper tip to any surface. Keep bottle tightly closed when not in use.

Please see full Prescribing Information, including Instructions for Use at www.CYSTADROPS.com.

 **Cystadrops**[®]
(cysteamine ophthalmic solution) 0.37%

3. Assist with dosing and instillation

To help your patients and caregivers instill the eyedrops successfully, assist them by:

- Reviewing the instructions on how to use CYSTADROPS
- Demonstrating yourself how to use the eyedrops
- Having them show you how they use the eyedrops

4. Discuss most common side effects

- The most common adverse reactions ($\geq 10\%$) are eye pain, vision blurred, eye irritation, ocular hyperaemia, instillation site discomfort, eye pruritus, lacrimation increased, and ocular deposits.
- Discussing side effects with patients and caregivers will help to prepare them for the possibility. Ask them to call your office if there are side effects and not to discontinue the eyedrops without first discussing the side effects with you.



- Tilt head back. Pull down lower eyelid with a finger.
- Gently squeeze the CYSTADROPS dropper to release 1 drop of CYSTADROPS into the space or "pocket" between the lower eyelid and the eye.



- Close the eye and press a finger into the corner of the eye by the nose, then gently massage the upper eyelid to spread the drop over the eye.
- Remove any medicine around the eye with a tissue.

Correct placement of the eyedrops:

- Results in less blinking and helps ensure that the eyedrop stays in the eye.
- Helps the drop coat the entire surface of the eye.

Important Safety Information (continued)

- Benign intracranial hypertension (or pseudotumor cerebri) associated with oral cysteamine or ophthalmic use of cysteamine (with concurrent oral cysteamine) has been reported, which has resolved with diuretic therapy.
- Contains benzalkonium chloride. Contact with soft contact lenses should be avoided. Remove contact lenses prior to application. Lenses may be reinserted 15 minutes following administration.
- The most common adverse reactions ($\geq 10\%$) are eye pain, vision blurred, eye irritation, ocular hyperaemia, instillation site discomfort, eye pruritus, lacrimation increased, and ocular deposits.
- **To report SUSPECTED ADVERSE REACTIONS, contact Recordati Rare Diseases Inc. at 1-888-575-8344, or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.**

Please see full Prescribing Information, including Instructions for Use at www.CYSTADROPS.com.



CYSTADROPS is a licensed trademark of Recordati Rare Diseases Inc.
Other trademarks, registered or otherwise, are the property of their respective owner(s).
© 2023 Recordati Rare Diseases Inc.
Recordati Rare Diseases Inc. • Bridgewater, NJ 08807
www.recordatirarediseases.com/us PP-CDPS-US-0159

 **Cystadrops**[®]
(cysteamine ophthalmic solution) 0.37%